

C13 to C15: Groups, rates and heat changes Knowledge Organiser

Lesson sequence

- Group 1
- Group 7
- Reactivity of halogens
- Group 0
- Rates of reaction
- Collision theory
- Core practical – rates of reaction (CP11)
- Catalysts
- Exothermic and endothermic reactions
- Explaining energy changes

1. Group 1

*Alkali metals	The name of the metals in group 1 – lithium, sodium, potassium and so on.
*Group 1 symbols	Li – lithium Na – sodium K – potassium
**Reaction of alkali metals with water	Metal + water → metal hydroxide + hydrogen E.g: sodium + water → sodium hydroxide + hydrogen $2\text{Na} + 2\text{H}_2\text{O} \rightarrow 2\text{NaOH} + \text{H}_2$
**Lithium and water	Lithium floats and bubble vigorously
**Sodium and water	Sodium melts into a ball and moves around the surface bubbling vigorously.
**Potassium and water	Potassium melts into a ball, catches fire (lilac) and moves around the surface bubbling vigorously.
*Group 1 reactivity	Reactivity increases as you move down the group.

**Explaining group 1 reactivity	When metals react they lose their outer electrons. Further down the group there are more shells of electrons so the outer electrons are less attracted to the nucleus and easier to remove.
--	---

2. Group 7

*Halogens	The names given to the non-metals in group 7 – fluorine, chlorine, bromine and iodine.
*Chlorine	Cl_2 . A pale green gas.
*Bromine	Br_2 . A red-brown liquid.
*Iodine	I_2 . A shiny purple-black solid.
**Reaction of halogens with metals	Halogen + metal → metal halide E.g: Bromine + sodium → sodium bromide $\text{Br}_2 + 2\text{Na} \rightarrow 2\text{NaBr}$
**Reaction of halogens with hydrogen	Halogen + hydrogen → hydrogen halide E.g: Chlorine + hydrogen → hydrogen chloride $\text{Cl}_2 + \text{H}_2 \rightarrow 2\text{HCl}$
*Hydrogen halides	Hydrogen halides dissolve in water to form acids, for example hydrogen chloride makes hydrochloric acid.
*Chlorine test	Chlorine gas turns damp blue litmus red then quickly bleaches it white.

3. Reactivity of halogens

*Group 7 reactivity	Reactivity increases as you go up the group.
**Explaining group 7 reactivity	When non-metals react they complete their outer shells. Further up the group the elements have fewer shells so the nucleus attracts electrons more strongly.

**Displacement reactions	Reactions in which a more reactive metal displaces a less reactive metal from a salt eg: $\text{copper sulfate} + \text{zinc} \rightarrow \text{zinc sulfate} + \text{copper}$ Does not work backwards as copper is less reactive than zinc.
**Displacement reactions of halogens	A more reactive halogen displaces a less reactive halide ion by taking its electrons. E.g: bromine + sodium iodide → iodine + sodium bromide
***Redox reactions of halogens	The more reactive halogen oxidises the less reactive halide by taking its electrons. The more reactive halogen is reduced. E.g: $\text{Br}_2 + 2\text{I}^- \rightarrow 2\text{Br}^- + \text{I}_2$

4. Group 0

*Noble gases	The name given to the non-metals in group 0 – helium, neon, argon, krypton and xenon.
*Melting point of noble gases	They are all gases at room temperature but the melting and boiling point increase down the group.
*Reactivity of group 0	The noble gases do not (easily) do any reactions – they are inert.
**Explaining reactivity of group 0	When elements react they try to complete their outer shells. Because group 0's outer shells are already complete, they do not react.
**Uses of noble gases	-Helium is used in airships because it is inert and has low density - Argon is used in fire extinguishers because it is inert and denser than air. - Neon is used in lighting because it glows red when electricity is passed through it.

5. Rates of reaction

*Rate of reaction	The rate at which reactants are used up or products are made.
*Reactants vs time graph	Starts high and curves downward, decreasing rapidly at first and then more gently. Steeper line = faster rate.
*Products vs time graph	Starts low and curves upwards, increasing rapidly at first and then more gently. Steeper line = faster rate.
**Measuring rates – reactions that produce gas	- Collect gas in a gas syringe and measure the volume every 30 secs. - Collect gas over water (up-turned measuring cylinder full of water) and measure volume every 30 secs. - Do reaction on a balance and record the change in mass every 30 secs.
**Measuring rates – reactions that go cloudy	Do the reaction in a beaker placed on piece of paper with a cross marked on it. Looking down through the beaker, time how it takes for the cross to disappear.

6. Collision theory


*Collision theory	States that for two particles to react they must: - Collide with each other - Collide with enough energy to react
*Activation energy	The minimum energy that two particles must have when they collide in order to react.
**Effect of concentration on rate	Increasing the concentration increases the rate because there are more particles so there are more collisions and more reactions.
**Effect of surface area on rate	Increasing the surface area (by decreasing particle sizes) increases the rate by exposing more particles to collisions leading to more collisions and more reactions.

**Effect of pressure on rate	Increasing the pressure increases the rate because particles are pushed closer together so they collide more often.
**Effect of temperature on rate	Increasing the temperature increases the rate because particles move faster so they collide more, and collide with more energy to a greater proportion of collisions lead to reactions.

7. Core practical – rates of reaction (CP11)


*CP11 – Aim	To explore the rate of two reactions by collecting gas and observing a colour change.
*CP11 – Gas collection – setup	Place a measuring cylinder full of water upside down in a basin of water. Place 5 g of marble chips in a conical flask with 40 cm ³ hydrochloric acid. Insert a bung with delivery tube and insert the delivery tube into the measuring cylinder.
*CP11 – Gas collection – measurements	Record the volume of gas collected every 15 seconds until it stops.
*CP11 – Gas collection – variations	Repeat with a different size of marble chips.
*CP11 – Gas collection – results	The amount of gas collected increases quickly at first and then more slowly. The smaller marble chips produce gas more quickly, but the same amount in total.
*CP11 – Colour change – setup	Draw a cross on a piece of paper and place a beaker on it. Measure out 50 cm ³ of sodium thiosulfate solution and 5 cm ³ of hydrochloric acid into two test tubes and leave to warm in a water bath at 30°C.

*CP11 – Colour change – run the experiment	Quickly pour both test tubes into the beaker, mix and start the stopwatch. Looking down through the beaker, stop when you can no longer see the cross.
*CP11 – Colour change – variations	Repeat with water baths set to 35°C, 40°C, 45°C and 50°C.
*CP11 – Colour change – results	The cross disappears most quickly at 50°C and least quickly at 30°C.




8. Catalyst

*Catalyst	A substance that speeds up a chemical reaction without being used up.
**Effect of catalysts on rate	Catalysts increase the rate of reaction by reducing the activation energy so that a greater proportion of collisions lead to reactions.
**Reaction profile	A graph that shows the changes in energy during a reaction. Starts with large 'hump' that represents the activation energy.
**Effect of catalysts on reaction profiles	The 'hump' representing the activation energy is smaller.
*Enzyme	A protein that works as a catalyst to speed up the reactions in our cells.
*Enzymes in alcohol production	Alcoholic drinks are produced using enzymes found in yeast which catalyse a reaction that turns glucose into ethanol.



9. Endothermic and exothermic reactions

*Exothermic reaction	A reaction that transfers energy to the surroundings (gets hotter).
*Exothermic reaction examples	- Neutralisation - Displacement - Combustion - Some precipitation - Respiration
*Endothermic reaction	A reaction that absorbs energy from the surroundings (gets colder)
*Endothermic reaction examples	- Dissolving (most) salts - Some precipitation - Photosynthesis
**Exothermic reaction profile	The reactants have more energy than the products, so their line on the graph is higher.
**Endothermic reaction profile	The reactants have less energy than the products, so their line on the graph is lower.
**Measuring energy changes	- Sit a polystyrene beaker inside a glass beaker (insulation) - Measure the starting temperature of the reactants. - Mix the reactants in the polystyrene beaker - Cover with lid fitted with a thermometer - Monitor and record the lowest temperature.




10. Explaining energy changes

**Chemical bonds in reactions	During chemical reactions, old chemical bonds are broken and new ones are formed.
**Breaking bonds	Breaking bonds absorbs energy, breaking stronger bonds absorbs more energy.
**Making bonds	Making bonds releases energy, making stronger bonds releases more energy.
**Energy changes and bond formation	The energy change in a reaction is the difference between the energy required to break the old bonds and the energy released by making the new ones.
**Exothermic reactions and bonds	Exothermic reactions break weaker bonds and make stronger ones.
**Endothermic reactions and bonds	Endothermic reactions break stronger bonds and make weaker ones.
***Bond strength	The energy required to break one mole of a particular covalent bond in kJ/mol.
***Calculating energy changes from bond strengths	Add up the total strength of old bonds broken and subtract the total strength of new bonds made. A negative answer is exothermic.

***Energy change example:

Hydrogen and chlorine react to form hydrogen chloride. The bond strengths are as follows: H-H = 436 kJ/mol, Cl-Cl = 240 kJ/mol, H-Cl = 428 kJ/mol. Calculate the energy change of the reaction



Bonds broken = 436 + 240 = 676
Bonds made = 2 x 428 = 856

Reaction energy = 676 – 856 = -180 kJ/mol, the reaction is exothermic because the answer is negative.

